Serial No.: 09/835,992

Conf. No.: 6680

- 3 -

Art Unit: 1642

In the Claims

Applicant has submitted a new complete claim set showing marked up claims insertions indicated by underlining and deletions indicated by strikeouts and/or double Applicant has submitted a new complete claim set showing marked up claims with bracketing.

1-12. (Cancelled)

- 13. (Currently amended) A method for determining gastric cancer in a patient, comprising, assaying a sample obtained from the patient for an antibody that specifically binds sterol carrier protein-X/sterol carrier protein-2, as a determination of the gastric cancer in said patient.
- 14. (Original) The method of claim 13, wherein the sterol carrier protein-X/sterol carrier protein-2 is encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NOs:19, 20, 21, and 22.
- 15. (Original) The method of claim 13, wherein the antibody is assayed using sterol carrier protein-X/sterol carrier protein-2 or an antigenic fragment thereof.

16-30. (Cancelled)

- 31. (Currently amended) A method for following the progress of a therapeutic regime designed to alleviate gastric cancer, comprising:
- (a) assaying a sample obtained from a subject at a first time to determine the level of a parameter selected from the group consisting of (i) a peptide derived from a sterol carrier protein X/sterol carrier protein-2, (ii) a cytolytic T cell specific for cells presenting said peptide, and (iii) an antibody which specifically binds sterol carrier protein-X/sterol carrier protein-2 to said peptide of said protein, at a first time period;

Serial No.: 09/835,992 - 4 - Art Unit: 1642

Conf. No.: 6680

(b) assaying the level of the antibody parameter selected in (a) obtained from the subject at a second period of time and comprising comparing it to the level determined in (a) as a determination of effect of said therapeutic regime.

32. (Previously presented) The method of claim 31, wherein the sterol carrier protein-X/sterol carrier protein-2 is encoded by a nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of SEQ ID NOs:19, 20, 21, and 22.

RULE 1.126 (New) The method of claim 32, wherein the antibody is assayed using sterol carrier protein-X/sterol carrier protein-2 or an antigenic fragment thereof.